

Time Asymmetric Quantum Physics

A. Bohm

Physics Department

The University of Texas at Austin

Austin, Texas 78712

Abstract

Mathematical and phenomenological arguments in favor of asymmetric time evolution of micro-physical states are presented.

1. Introduction

Standard quantum mechanics in Hilbert space $\mathcal{H}$ is a time symmetric theory with a time symmetric dynamical (differential) equation and time symmetric boundary conditions. This is in contrast to many time asymmetric phenomena observed in classical and also in quantum physics. Of the latter we want to discuss in this article two examples, the decay of a quasi-stable particle [1] and the expanding universe as a whole when considered as a closed (without extrinsic influences [2]) quantum systems [3].

In classical physics solutions of time-symmetric dynamical equations with time asymmetric boundary conditions come in pairs, e.g., big bang – big crunch in general relativity or retarded – advanced in electromagnetism. With the choice of the boundary condition, one out of the two time-asymmetric solutions is selected. The Hilbert space theory of quantum mechanics does not allow such time-asymmetric formulations. In the Hilbert space formulation of quantum mechanics the space-time transformations (e.g., Galilean transformations, Poincaré transformations) are described by a unitary group representation in the Hilbert space $\mathcal{H}$. Thus the time evolution is unitary and reversible, and it is given by $U^\dagger(t) = \exp(-iHt)$, $-\infty < t < \infty$. This is the consequence of a series of mathematical theorems which follow from the mathematical properties - specifically the topological completeness - of the Hilbert space; they are listed in the Appendix A. These theorems in particular exclude the existence of non-zero probabilities which are zero before a given finite time t_0 ($t_0 \neq -\infty$), which is the time at which the quasi-stable particle had been produced or the time of the big bang in the two examples of this article. The decay of resonances and the quantum theory of our universe can therefore not be described consistently in the mathematical theory using the Hilbert space.

Disregarding Hilbert space mathematics, in scattering theory one arrived in a heuristic way at a pair of time asymmetric boundary conditions by choosing in- and out-plane wave “states” $|E^+\rangle$ and $|E^-\rangle$ which have their origin in the $\epsilon = +0$ and $\epsilon = -0$ of the Lippmann-Schwinger equation [4], cf. Appendix B. Still, the widespread opinion remained

that asymmetric or irreversible time evolution of closed quantum mechanical systems is impossible.

It could have been that historically the analogy to classical mechanics was the origin of this belief, though the time evolution for the Schrödinger equation could have as well been discussed in analogy to the electromagnetic waves, and for those the radiation arrow of time was well accepted (and by some even considered as fundamental [5]). However, the reversibility of the Hamiltonian generated time evolution in von Neumann's [6] Hilbert space quantum theory must have been a decisive factor for the longevity of this belief.

Already the Dirac [7] kets $|E\rangle$, $0 \leq E < \infty$, are not elements of the Hilbert space but generalized eigenvectors and required the extension of the Hilbert space $\mathcal{H}$ to the Rigged Hilbert Space $\Phi \subset \mathcal{H} \subset \Phi^\times$ [8], where Φ is a linear scalar product space of well-behaved vectors $\phi \in \Phi$ (represented by smooth *etc.* functions $\langle E|\phi\rangle$) and $\Phi^\times \ni |E\rangle$ is the space of its antilinear functionals. In the S -matrix element [cf. Appendix B]

$$(1.1) \quad (\psi^{\text{out}}, S\phi^{\text{in}}) = (\psi^-, \phi^+) = \sum_{bb'} \int_0^\infty dE \langle \psi^- | b, E^- \rangle \langle b | S(E) | b' \rangle \langle +b', E | \phi^+ \rangle$$

appear the Dirac “scattering states” $|E^\pm\rangle$ which are obtained from $|E\rangle$ by the Lippmann-Schwinger equation. In order to analytically continue to the resonance pole $z_R = E_R - i\Gamma/2$ of the S -matrix $\langle b | S(z) | b' \rangle$ the set of in-states $\{\phi^+\} \equiv \Phi_-$ and out-states $\{\psi^-\} \equiv \Phi_+$ must additionally have some analyticity property. In order to get a Breit-Wigner energy distribution for the pole term we postulate that the energy wave functions $\langle -E | \psi^- \rangle$ and $\langle +E | \phi^+ \rangle$ are well-behaved Hardy class functions of the upper and lower half-plane in the second sheet of the energy surface of the S -matrix.

The analytically continued Dirac kets $|E^-\rangle \in \Phi_+^\times$ of the Lippmann-Schwinger equations become – using the Cauchy formula – at the resonance pole $z_R = E_R - i\Gamma/2$ the Gamow kets $|z_R^-\rangle \in \Phi_+^\times$. The time asymmetric semigroup evolution of these Gamow kets

$$(1.2) \quad e^{-iH^\times t} \Big|_{t \geq 0} |z_R^-\rangle = e^{-iE_R t} e^{-\Gamma/2 t} |z_R^-\rangle, \quad \text{for } t \geq 0 \text{ only,}$$

is then derived as a mathematical consequence of the structure of the Rigged Hilbert Space $\Phi_+^\times \supset \mathcal{H} \supset \Phi_+$ of Hardy class [9] (in the same way as the time symmetric unitary group evolution given by $e^{-iH^\dagger t}$ $-\infty < t < +\infty$, is a mathematical consequence of the Hilbert space structure).

Thus asymmetric time evolution would be a natural property of quantum mechanical states represented by the vector $|z_R^-\rangle$ and other elements of the space $\Phi_+^\times$. In this article we want to discuss the phenomenological evidence for such states and the experimental conditions and phenomenological reason for the asymmetric time evolution.

In section 2 we review the basic concepts of quantum physics in a way that shows which mathematical properties are important for quantum mechanical calculations and which are idealizations and not directly obtainable from experimental data. We also argue that experimental observations involve a time asymmetry, the preparation $\Rightarrow$ registration arrow of time. We then give two examples of quantum mechanical states with asymmetric time evolution, the quasi-stable particle and the universe considered as a closed quantum system, and discuss their common features. In section 3 we provide the mathematical theory for time asymmetric quantum mechanics and give some of its result. In section 4 we discuss an example of a state with an arrow of time prepared in a laboratory experiment; we compare the concept of its preparation time with the initial time for the state of the quantum universe.

2) Calculational Methods, Mathematical Idealizations and Experimental Observations

In quantum theory one has states and observables. States are described by density or statistical operators and conventionally denoted by ρ or W ; for pure states vectors ϕ are used. Observables are described by operators $A(= A^\dagger)$, Λ , $P(= P^2)$, but we will also use vectors ψ to describe a state P if $P = |\psi\rangle\langle\psi|$.

The vectors ϕ, ψ are elements of a vector space Φ with a scalar product, denoted $(\cdot, \cdot)$ or $\langle \cdot | \cdot \rangle$. The operators A, Λ , are elements of the algebra of linear operators $\mathcal{A}$ in Φ . The linear space Φ , though often called a Hilbert space, is mostly treated like a pre-Hilbert space, i.e., without a topology (or without a definition of convergence) and it is not topologically complete. If we want to emphasize that Φ has no topology we denote it by Φ_{alg} .

Each “kind” of quantum physical system is associated to a space Φ .

In experiments, the state W (or the pure (idealized) state ϕ) is prepared by a preparation apparatus and the observable A (or the idealized observable ψ) is registered by a registration apparatus (e.g., a detector). The fundamental aspect of the new theory presented here is to clearly distinguish between states (e.g., in-states ϕ^+ of a scattering experiment) and observables (e.g., detected out-states ψ^- of a scattering experiment), cf. Appendix B.

The measured (or registered) quantities are ratios of (usually) large numbers, the detector counts. They are interpreted as probabilities, e.g., as the probability to measure the observable Λ in the state W at the time t , which is denoted by $\mathcal{P}_W(\Lambda(t))$.

The probabilities are calculated in theory as the scalar product or, in the general case, as the trace. This is shown in relations (2.1a) and (2.1b), below where $\approx$ indicates the equality between the experimental and the theoretical quantities and $\equiv$ is the mathematical definition of the theoretical probabilities in terms of the quantities of the space Φ (which

is not yet completely defined):

$$(2.1a) \quad N_i/N \approx \mathcal{P}_\phi(P) \equiv |\langle \psi | \phi \rangle|^2$$

$$(2.1b) \quad N(t)/N \approx \mathcal{P}_W(\Lambda(t)) \equiv \text{Tr}(\Lambda(t)W_0) = \text{Tr}(\Lambda_0 W(t))$$

The parameter t in (2.1b) is the continuous time parameter and the observable Λ , or the state W , are “continuous” functions of time (with $W_0 = W(t=0)$). Thus, $\mathcal{P}_W(\Lambda(t))$ is thought of as a continuous function of t . But $N(t)$ is the number of counts in the time interval between $t=0$ and t , which is an integer. Thus the right hand side of $\approx$ in (2.1b) changes continuously in t , but the left hand side can only change in steps of rational numbers. This shows that the continuity of $\mathcal{P}_W(\Lambda(t))$ or of $|\langle \psi(t) | \phi \rangle|^2 = |\langle \psi | \phi(t) \rangle|^2$ as a function of t , and similar topological questions, are not directly experimentally testable.

For more general observables A , which are expressed in terms of the orthogonal projection operators P_i ($P_i P_j = \delta_{ij} P_j$) as

$$(2.2) \quad A = \sum_{i=1}^{\infty} a_i P_i,$$

where a_i are the eigenvalues of A , the probabilities are measured as the average value $\sum_{i=1}^{\text{finite}} a_i \frac{N_i}{N}$. Here the sum is finite since an experiment can give only a finite number of data. In the comparison between theory and experiment this finite sum is represented by the infinite sum obtained from (2.2), thus

$$(2.3) \quad \sum_{i=1}^{\text{finite}} a_i \frac{N_i}{N} \approx \mathcal{P}(A) = \sum_{i=1}^{\infty} a_i \mathcal{P}(P_i).$$

This also shows that the meaning of such topological notions as the convergence of infinite sequences (of e.g., partial sums of the right hand side of (2.2) and (2.3)) cannot be established directly from the experimental data on the left hand side of (2.3), which provides only a finite sequence. Thus the definition of convergence of infinite sequences in Φ , i.e.

the topology of the space Φ , is a mathematical idealization. If one wants a complete mathematical theory one needs to make this mathematical idealization and choose a topology for the space Φ . Usually, for many practical calculations in physics, one does not worry about the completeness and uses instead some calculational rules.

To obtain the rules for calculating the trace and the scalar product on the right hand side of (2.1) one starts with a basis vector decomposition for the state vector $\phi \in \Phi$ using a discrete set of eigenvectors $|i\rangle = |\lambda_i\rangle$ of an observable (often the Hamiltonian) with eigenvalues λ_i .

$$(2.4) \quad \phi = \sum |i\rangle \langle i|\phi\rangle$$

Often, following Dirac [7], one uses a continuous set of eigenvectors $|\lambda\rangle$ (Dirac kets) and writes:

$$(2.5) \quad \phi = \int d\lambda |\lambda\rangle \langle \lambda|\phi\rangle$$

The trace, scalar product, etc., are then calculated as

$$(2.6a) \quad \text{Tr}(\Lambda W) = \sum_i \langle i|\Lambda W|i\rangle \quad \text{or, (2.6b)} \quad \text{Tr}(\Lambda W) = \int d\lambda \langle \lambda|\Lambda W|\lambda\rangle$$

$$(2.7a) \quad |\langle \psi|\phi\rangle|^2 = \left| \sum_{i=1}^{\infty} \langle \psi|i\rangle \langle i|\phi\rangle \right|^2 \quad \text{or, (2.7b)} \quad |\langle \psi|\phi\rangle|^2 = \left| \int d\lambda \langle \psi|\lambda\rangle \langle \lambda|\phi\rangle \right|^2$$

In practical calculations the convergence of infinite sums and the meaning of integration (Lebesgue versus Riemann) are usually not considered. Often one truncates to finite (e.g., two) dimensions such that of the sums in (2.6a) and (2.7a) one retains only a finite number of terms. If one has a complete mathematical theory one can define the meaning of the infinite sums in (2.6a) (2.7a) and the meaning of the integrals in (2.6b), (2.7b) and prove (2.4) and (2.5). For instance one can choose for Φ the Hilbert space $\mathcal{H}$, in which case (2.4) but not (2.5) can be proven. Or one can choose for Φ a complete space with some locally convex, nuclear topology and its space of continuous functionals $\Phi^\times$ to obtain a Gelfand

triplet $\Phi \subset \mathcal{H} \subset \Phi^\times$. Then the kets are $|\lambda\rangle \in \Phi^\times$ and one can prove the Dirac basis vector expansion (2.5) as the Nuclear Spectral Theorem.

Time evolution, i.e., the dynamics of a quantum physical system, is given by the Hamilton operator H of the system. (H is always assumed to be (essentially) self-adjoint, $\bar{H} = H^\dagger$, and semibounded). The dynamical equation is the von Neumann or Schroedinger equation:

$$(2.8) \quad \frac{\partial W(t)}{\partial t} = \frac{i}{\hbar} [H, W(t)]; \quad i\hbar \frac{\partial \phi(t)}{\partial t} = H^\dagger \phi(t)$$

$$\phi(t=0) = \phi_0$$

Equivalently, one gives the time evolution in the Heisenberg picture by

$$\frac{\partial \Lambda(t)}{\partial t} = -\frac{i}{\hbar} [H, \Lambda(t)]; \quad i\hbar \frac{\partial \psi(t)}{\partial t} = -H\psi(t)$$

$$\psi(t=0) = \psi_0$$

In a time symmetric theory, that means if one uses for the time symmetric differential equation (2.8) *also* time symmetric boundary conditions, then, one obtains the following solutions of (2.8):

$$(2.9) \quad W(t) = e^{-iHt} W_0 e^{iHt}, \text{ where } -\infty < t < \infty,$$

$$(2.10) \quad \phi(t) = U^\dagger(t) \phi_0 = e^{-iHt} \phi_0, \quad -\infty < t < \infty$$

or, in the Heisenberg picture,

$$(2.11) \quad \Lambda(t) = e^{iHt} \Lambda_0 e^{-iHt}, \quad \text{where } -\infty < t < \infty,$$

Here $\Lambda_0 \equiv \Lambda(t=0)$, $W_0 \equiv W(t=0)$.

On the other hand if one just starts with the differential equations (2.8) and postulates the Hilbert space topology, $\phi(t) \in \mathcal{H}$, then the above unitary group evolution is

the only possible solution of the dynamical equations (this follows from some theorems of Gleason and Stone (Appendix A)). This means time asymmetric boundary conditions which could result in an irreversible time evolution are not mathematically allowed in a quantum theory in the Hilbert space $\mathcal{H}$. The assumption $\phi(t) \in \mathcal{H}$ always leads to the time evolution (2.10) given by the unitary group $U(t)$ which has always an inverse $U(-t)$. Inserting (2.9), (2.10) or (2.11) into the right hand side of (2.1), the probability $\mathcal{P}(t) = \text{Tr}(\Lambda W(t))$ can be calculated at any time $t_0 + t$ or $t_0 - t$.

In contrast to the results calculated with (2.9), the probabilities $\mathcal{P}(t)$ cannot be observed at any arbitrary positive or negative time t . The reason is the following:

A state needs to be prepared before an observable can be measured, or registered in it.

We call this truism the preparation $\Rightarrow$ registration arrow of time [18]; it is an expression of causality. Let $t_0(=0)$ be the time at which the state has been prepared. Then, $\mathcal{P}(\Lambda(t))$ is measured as the ratio of detector counts

$$(2.12a) \quad \mathcal{P}_W^{\text{exp}}(\Lambda(t)) \approx \frac{N(t)}{N},$$

$$(2.12b) \quad \text{for } t > t_0 = 0,$$

If there are some detector counts before $t = t_0$, they are discounted as noise because the experimental probabilities

$$(2.13) \quad \text{can not fulfill } \mathcal{P}_W^{\text{exp}}(\Lambda(t)) \not\approx 0, \quad \text{for } t < t_0 = 0,$$

Though in the Hilbert space theory $\mathcal{P}_W(\Lambda(t)) = \mathcal{P}_{W(t)}(\Lambda)$ can be calculated at positive or negative values of $t - t_0$ using unitary group evolution (2.9), an experimental meaning can be given to $\mathcal{P}_W(\Lambda(t))$ only for $t > t_0$.

In some cases (e.g., stationary states, cyclic evolutions), it should not matter at what time $\mathcal{P}_{W(t)}(\Lambda)$ is calculated because one can extrapolate to negative values of t .

The physical question is: Are there quantum physical states in nature that evolve only into the positive direction of time, $t > t_0$, and for which one therefore cannot extrapolate to negative values of $t - t_0$? If there are such states, pure states or mixtures, they

cannot be described by the standard Hilbert space quantum theory, because of the unitary group time evolution (2.9) and (2.10), which is a mathematical consequence of the specific (topological, not algebraic) structure of the Hilbert space.

Two prominent examples of states with an asymmetric time evolution, $t > t_0$, are the decaying states (in all areas of physics, relativistic or non-relativistic) and our universe as a whole, considered as a quantum physical system.

1.) Decaying states and resonances are often thought of as something complicated, because in the Hilbert space there does not exist a vector that can describe them in the same way as stable states are described by energy eigenvectors. However, empirically, quasi-stable particles are not qualitatively different from stable particles; they differ only quantitatively by a non-zero value of the width Γ . Stability or the value of lifetime is not taken as a criterion of elementarity, at least not by the practitioners [10]. A particle decays if it can and it remains stable if selection rules for some quantum numbers prevent it from decaying. Therefore, stable and quasi-stable states should be described on the same footing, e.g., define both by a pole of the S-matrix at the position $z_R = E_R - i\Gamma/2$, or/and as a generalized eigenvector with eigenvalue z_R (with $\Gamma = 0$ for stable particles). Since the latter is not possible in the Hilbert space, one devises “Effective Theories” in order to obtain a state vector description of quasi-stable states.

Phenomenological effective theories have been enormously successful. They describe resonances in a finite dimensional space as eigenvectors of the “effective Hamiltonian” with complex eigenvalue $(E_R - i\Gamma/2)$, where E_R =resonance energy, $\hbar/\Gamma$ =life time, and their time evolution is given by the exponential law. The common feature of these approximate methods is the omission of a continuous sum; the infinite dimensional theory is truncated to a finite (e.g., two) dimensional effective theory. Examples of this approach are: The approximate method of Weisskopf and Wigner and of Heitler for atomic decaying states [11]; the Lee-Oehme-Yang effective two dimensional theory for the neutral Kaon system [12]; and many more finite dimensional models with non-Hermitian diagonalizable Hamiltonian

matrices in nuclear physics [14]. Also non-diagonalizable finite dimensional Hamiltonians were discussed [13]. In the Hilbert space framework “there does not exist . . . , a rigorous theory to which these methods can be considered as approximations” [15].

The decay of a quantum physical system, e.g., the transition of an excited state of a molecule into its ground state or the decay of an elementary particle [16] is a profoundly irreversible process. Therefore we should like to introduce state vector $|F\rangle$, $|\psi^G\rangle = |E_R - i\Gamma/2\rangle$ or state operators $W^G(t) = |F\rangle\langle F|$, for which the time evolution is asymmetric and for which the theoretical probabilities $\text{Tr}(\Lambda W^G(t))$ can be calculated for $t > t_0 = 0$ only.

This means we have to generalize the unitary group evolution (2.9), (2.10) with $-\infty < t < \infty$ to a semigroup evolution with $0 \leq t < \infty$. This is accomplished by seeking solutions of the time symmetric dynamical equations (2.8) with time *asymmetric* boundary conditions. Since in Hilbert space quantum mechanics semigroup evolution is not possible, we seek a semigroup solution $F(t)$ to the quantum mechanical Cauchy problem (2.8) with Hamiltonian $H_+^\times$ where $F(t)$ is an element of a larger space in which $\mathcal{H}$ is dense and which we denote by $\Phi_+^\times \supset \mathcal{H}$, i.e., the Hamiltonian $H_+^\times$ is the uniquely defined extension of the Hilbert space Hamiltonian $H^\dagger$ to this space $\Phi_+^\times$. Thus the dynamical equation (2.8) is:

$$(2.14) \quad i\hbar \frac{\partial F(t)}{\partial t} = H_+^\times F(t)$$

with the initial data $F(t=0) = F_0^- \in \Phi_+^\times$, and the solution is given by the semigroup¹

$$(2.15a) \quad F(t) = U_+^\times(t) F_0^- = e_+^{-iH^\times t} F_0^-$$

$$(2.15b) \quad \text{for } t \geq 0 \text{ only.}$$

¹ This semigroup is generated by the Hamiltonian H :

$$|F(t)\rangle\langle F(t)| = e_+^{-iH^\times t} |F(0)\rangle\langle F(0)| e_+^{iHt}$$

It is *not* the semigroup of quantum statistical mechanics of open systems generated by a

If we use the quantum mechanical state operators with semigroup time evolution,

$$(2.16) \quad W^G(t) \equiv |F(t)\rangle\langle F(t)| = e^{-iH^\times t} W^G(t_0) e^{iHt}, \quad t \geq 0,$$

to calculate the quantum mechanical probabilities, then for these calculated probabilities we obtain

$$(2.17a) \quad P_{W^G(t)}(\Lambda_0) = \text{Tr}(\Lambda(t_0)W^G(t)) = \text{Tr}(\Lambda(t)W^G(t_0))$$

$$(2.17b) \quad t \geq t_0 = 0.$$

This means that they fulfill the same conditions as the experimental probabilities (2.12a), (2.12b) and (2.13).

In particular the probabilities are not defined unless the preparation $\Rightarrow$ registration arrow of time (2.17b) is fulfilled, because the time evolution

$$(2.18a) \quad W^G(t) = e^{-iH^\times t} W_0^G e^{iHt} \quad \text{or} \quad (2.18a') \quad \Lambda(t) = e^{iHt} \Lambda_0 e^{-iH^\times t}$$

Liouvillian L , i.e., this is not the irreversible time evolution of open systems under external influences [17]. For the state ρ of such open systems, one has in place of (2.8)

$$\frac{\partial \rho(t)}{\partial t} = L\rho(t) = -i[H, \rho(t)] + \mathcal{I}\rho(t),$$

where H is the Hamiltonian of the open system and $\mathcal{I}$ is the interaction of the external reservoir upon the system, e.g.,

$$\mathcal{I} = \sum_{\alpha=1,2,..} ([V_\alpha \rho(t), V_\alpha^\dagger] + [V_\alpha, \rho(t) V_\alpha^\dagger])$$

The time evolution semigroup for open systems is

$$\rho(t) = \Lambda(t)\rho(0), \quad \text{where } \Lambda(t) = e^{Lt}, \quad t \geq 0$$

$\rho(t)$ describes the state of an *open* system acted upon by an external reservoir (environment, measurement apparatus, quantum reservoir, etc.)

is a semigroup evolution and only defined for

$$(2.18b) \quad t > t_0 = 0.$$

The physical meaning of the initial time t_0 for a decaying system in the state W^G will be discussed in section 4 below. Mathematically, it is given by the initial time $t = 0$ of the Cauchy problem (2.14).

This semigroup-arrow of time (2.15b), (2.17b), (2.18b) is the formulation in the mathematical theory of the experimental preparation $\Rightarrow$ registration arrow of time (2.12).²

2.) The universe, when considered as a quantum physical system, must also be in a state ρ (a pure state $\rho = |\phi\rangle\langle\phi|$, or a mixture) with asymmetric time evolution [19]. Its arrow of time must be identical with the traditional cosmological arrow of time and the time $t = t_0 = 0$, at which the initial state of the universe ρ has been prepared, is the time of the big bang.

The general quantum mechanical (a priori) probabilities predicted for the observable represented by the projection operator $P_{\alpha_1}^1(t_1)$ (“yes-no observations”) are according to (2.17),

$$(2.19a) \quad \mathcal{P}(\alpha_1, t_1) \equiv \mathcal{P}_\rho(P_{\alpha_1}^1(t_1)) = \text{Tr}(P_{\alpha_1}^1(t_1)\rho) = \text{Tr}(P_{\alpha_1}^1(t_1)\rho P_{\alpha_1}^1(t_1))$$

$$(2.19b) \quad \text{for } t_1 > t_0 = 0 \text{ only}$$

The time ordering (2.19b) is the same as the semigroup arrow of time (2.17b) in the quantum mechanics of measured systems. Applied to experiments performed on quantum

² Since the semigroup time evolution (2.15) or (2.18) is not possible in the Hilbert space, i.e., $F_0^- \notin \mathcal{H}$, people who wanted to retain the standard Hilbert space theory but were aware of the quantum mechanical preparation $\Rightarrow$ registration arrow of time had to extrapolate (2.18) to negative times, therewith eliminating the experimental preparation $\Rightarrow$ registration arrow of time and causality from the mathematical theory [18].

systems in the laboratory it leads to the preparation $\implies$ registration arrow of time (2.12b). Like in the quantum mechanics of measured systems, (2.19b) is an expression of causality.

The quantum mechanical probabilities (2.19) of projection operators $P_{\alpha_i}^i(t_i)$ can be generalized to probabilities of histories [3, 20].

A history is a time ordered product of different projection operators (labeled by α_i) for different observables (labeled by i):

$$(2.20) \quad C_\alpha = P_{\alpha_1}^1(t_1) \dots P_{\alpha_i}^i(t_i) \dots P_{\alpha_n}^n(t_n); \quad t_n > t_{n-1} > \dots > t_2 > t_1.$$

with

$$(2.21a) \quad P_{\alpha_i}^i(t_i) = e^{iH(t_i-t_{i-1})} P_{\alpha_i}^i(t_{i-1}) e^{-iH(t_i-t_{i-1})},$$

$$(2.21b) \quad t_i - t_{i-1} > 0$$

This definition of histories is suggested by the following considerations:

Let $P_{\alpha_i}^i$ be the α_i -th projector of (what we denote as) the i -th observable $A^i = \sum_\alpha a_\alpha^i P_\alpha^i$ $i = 1, 2, 3, \dots$. Then, starting with the operator $\rho = \rho(t_0)$ of (2.19a), one can define a sequence of effective density operators $\rho^{\text{eff}}(t_1), \dots, \rho^{\text{eff}}(t_{n-1})$, and one can predict a sequence of probabilities $\mathcal{P}(\alpha_2 t_2; \alpha_1 t_1), \mathcal{P}(\alpha_3 t_3; \alpha_2 t_2; \alpha_1 t_1) \dots \mathcal{P}(\alpha_n t_n; \dots \alpha_1 t_1)$. These density operators and probabilities are listed below:

$$(2.22a) \quad \rho^{\text{eff}}(t_1) = \frac{P_{\alpha_1}^1(t_1) \rho(t_0) P_{\alpha_1}^1(t_1)}{\text{Tr}(P_{\alpha_1}^1(t_1) \rho(t_0) P_{\alpha_1}^1(t_1))} = N_1 P_{\alpha_1}^1(t_1) \rho(t_0) P_{\alpha_1}^1(t_1)$$

$$(2.22b) \quad \text{for } t_1 > t_0 \text{ only;}$$

(the second equality in (2.22a) defines the normalization factor N_1)

and

$$(2.23a) \quad \mathcal{P}(\alpha_2 t_2; \alpha_1 t_1) = N_1 \text{Tr}(P_{\alpha_2}^2(t_2) \rho^{\text{eff}}(t_1) P_{\alpha_2}^2(t_2))$$

$$(2.23b) \quad \text{for } t_2 > t_1 \text{ only.}$$

Continuing in this way for $n = 3, 4, \dots$,

$$\rho^{\text{eff}}(t_{n-1}) = \frac{P_{\alpha_{n-1}}^{n-1}(t_{n-1})\rho^{\text{eff}}(t_{n-2})P_{\alpha_{n-1}}^{n-1}(t_{n-1})}{\text{Tr}(P_{\alpha_{n-1}}^{n-1}(t_{n-1})\rho^{\text{eff}}(t_{n-2})P_{\alpha_{n-1}}^{n-1}(t_{n-1}))}$$

$$(2.24a) \quad = N_{n-1}P_{\alpha_{n-1}}^{n-1}(t_{n-1})\rho^{\text{eff}}(t_{n-2})P_{\alpha_{n-1}}^{n-1}(t_{n-1});$$

$$= \frac{P_{\alpha_{n-1}}^{n-1}(t_{n-1}) \cdots P_{\alpha_1}^1(t_1)\rho(t_0)P_{\alpha_1}^1(t_1) \cdots P_{\alpha_{n-1}}^{n-1}(t_{n-1})}{\text{Tr}(P_{\alpha_{n-1}}^{n-1}(t_{n-1}) \cdots P_{\alpha_1}^1(t_1)\rho(t_0)P_{\alpha_1}^1(t_1) \cdots P_{\alpha_{n-1}}^{n-1}(t_{n-1}))}$$

$$(2.24b) \quad t_{n-1} > t_{n-2} > \cdots > t_1 > t_0$$

and

$$(2.25a) \quad \mathcal{P}(\alpha_n t_n; \cdots \alpha_1 t_1) = \frac{\text{Tr}(P_{\alpha_n}^n(t_n)\rho^{\text{eff}}(t_{n-1}))}{\text{Tr}(P_{\alpha_{n-1}}^{n-1}(t_{n-1})\rho^{\text{eff}}(t_{n-2})P_{\alpha_{n-1}}^{n-1}(t_{n-1}))};$$

$$(2.25b) \quad t_n > t_{n-1}$$

or

$$(2.26a) \quad \mathcal{P}(\alpha_n, t_n; \dots; \alpha_1, t_1) = N_{n-1} \text{Tr}(P_{\alpha_n}^n(t_n) \cdots P_{\alpha_1}^1(t_1)\rho(t_0)P_{\alpha_1}^1(t_1) \cdots P_{\alpha_n}^n(t_n));$$

$$(2.26b) \quad \text{for } t_n > \dots > t_0 \text{ only}$$

The time ordering or arrow of time (2.23b) ... (2.26b) is again the same as the semi-group arrow of time (2.18b), and (2.21) is the same as the semigroup evolution (2.18') (in the Heisenberg picture) for the observable Λ in the quantum theory of measured systems.

The probability (2.25a), (2.26a) is the probability of the history defined in (2.20)

$$(2.27) \quad \mathcal{P}(\alpha_n t_n \dots \alpha_1 t_1) = N_n \text{Tr}(C_\alpha \rho(t_0) C_\alpha)$$

One can consider alternative projection operators

$$(2.28a) \quad C'_\alpha = P_{\alpha'_1}^{1'}(t_1)P_{\alpha'_2}^{2'}(t_2) \cdots P_{\alpha'_n}^{n'}(t_n)$$

but a physical meaning can only be given to these products for the time ordering

$$(2.28b) \quad t_n > t_{n-1} \dots > t_1.$$

This time ordering, identical with the time ordering (2.22b) ... (2.24b), is a calculational consequence of the restriction (2.21b) postulated [3, 19] for the time evolution of the projectors. The restricted time evolution (2.21) is a semigroup evolution generated by the Hamiltonian of the closed quantum system. Obviously the semigroup (2.21), (2.18) and (2.16) is the same semigroup applied to different observables, $P_{\alpha_i}^i$ and Λ respectively of different quantum systems, namely the quantum universe and the quasi-stable particle. The semigroup character of the time evolution (2.18') — or of (2.18) in the Schroedinger picture — was inferred from restrictions imposed by observational limitations in a laboratory experiment with quantum systems, namely from the preparation $\Rightarrow$ registration arrow of time. The semigroup character of the time evolution (2.21) and the time ordering (2.28b) were postulated for the quantum universe because of the special initial state associated to the big bang [19]. From the way the time ordering appears in the probabilities for the laboratory experiments (2.17) and in the probabilities of the histories (2.19) ... (2.25), it is clear that both time orderings express the same arrow of time. If our universe is a closed quantum system as suggested by [3], the semigroup arrow for the resonances is subsumed under the cosmological arrow of time, or vice versa. This arrow of time “may *not* be attributed to the thermodynamic arrow of an external measuring apparatus (for the laboratory experiment) or larger universe” (for the quantum universe). It is a “fundamental quantum mechanical distinction between the past and future” [3].

As mentioned above, a semigroup evolution that could give a theoretical description of this arrow of time is impossible in the standard Hilbert space quantum mechanics. Therefore, in order to make the semigroup postulate (2.21) possible and to allow for a semigroup solution (2.15) of the quantum mechanical Cauchy problem, one must develop a new mathematics. We shall present the mathematics that is capable of a time asymmetric quantum theory in the following section.

3. A Mathematical Theory for Time Asymmetric Quantum Physics

Our empirical consideration in section 2 has led us to the postulate of a time evolution semigroup (2.21) or (2.18). Here we want to discuss a mathematical theory of quantum physics for which a semigroup evolution exists.

In a linear space with a scalar product Φ_{alg} , which we need for the calculational rules of quantum mechanics, the simplest modification that allows Hamiltonian generated semigroups is to choose instead of the Hilbert space topology a locally convex topology. If one also wants the Dirac formalism (i.e., kets, the basis vector expansion (2.5) etc.), then one has to choose a Rigged Hilbert Space (RHS) or Gelfand triplet.

$$(3.1) \quad \Phi \subset \mathcal{H} \subset \Phi^\times$$

The triplet of spaces in a Rigged Hilbert Space $\Phi \subset \mathcal{H} \subset \Phi^\times$ results from three different topological completions of the same algebraic (pre-Hilbert) space Φ_{alg} of section 2, [21]. Completion means adjoining to Φ_{alg} the (limit elements of) convergent (Cauchy) sequences with respect to a topology. The completion of Φ_{alg} with respect to the norm $\|\varphi\| = \sqrt{(\varphi, \varphi)}$, $\varphi \in \Phi_{\text{alg}}$ is the Hilbert space $\mathcal{H}$. The topology or meaning of convergence defined by the norms we denote by $\mathcal{T}_{\mathcal{H}}$. The completion of Φ_{alg} with respect to a finer locally convex, nuclear topology, which we denote by $\mathcal{T}_{\Phi}$ (and which is usually given by a countable number of norms [21]), is denoted by Φ . Then one has $\Phi_{\text{alg}} \subset \Phi \subset \mathcal{H}$ (because Φ and $\mathcal{H}$ contain all elements of Φ_{alg} plus the limit elements of Cauchy sequences in Φ_{alg}), and $\Phi \subset \mathcal{H}$ holds because $\mathcal{T}_{\Phi}$ is chosen to be finer or stronger than $\mathcal{T}_{\mathcal{H}}$ and there are consequently more $\mathcal{T}_{\mathcal{H}}$ Cauchy sequences than $\mathcal{T}_{\Phi}$ Cauchy sequences. We also consider the space of $\mathcal{T}_{\mathcal{H}}$ -continuous and of $\mathcal{T}_{\Phi}$ -continuous functionals. $\mathcal{H}^\times$ is the space of $\mathcal{T}_{\mathcal{H}}$ continuous antilinear functionals ψ on the space $\mathcal{H}$: $\psi : \phi \in \mathcal{H} \rightarrow \psi(\phi) \in \mathcal{C}$, and $\mathcal{H} = \mathcal{H}^\times$, $\psi(\phi) = (\phi, \psi)$, by a mathematical theorem. $\Phi^\times$ is the space of $\mathcal{T}_{\Phi}$ -continuous, antilinear functionals F on the space Φ : $F : \phi \in \Phi \rightarrow F(\phi) \equiv \langle \phi | F \rangle \in \mathcal{C}$. One has $\mathcal{H}^\times \subset \Phi^\times$ and the bra-ket $\langle | \rangle$ becomes an extension of the scalar product. Thus one obtains the Gel'fand triplet (3.1).

Dirac kets are elements of $\Phi^\times$, but there are also other $|F\rangle \in \Phi^\times$ besides the Dirac kets. Dirac's algebra of observables is an algebra of continuous operators in Φ (observables cannot be continuous operators in $\mathcal{H}$).

For a $\mathcal{T}_\Phi$ -continuous linear operator A , its conjugate operator $A^\times$ is defined by

$$(3.2) \quad \langle A\phi|F\rangle = \langle \phi|A^\times|F\rangle, \quad \forall \phi \in \Phi \text{ and } \forall F \in \Phi^\times$$

$A^\times$ is a continuous operator in $\Phi^\times$. Then for each observable A , one has a triplet of operators

$$(3.3) \quad A^\dagger|_\Phi \subset A^\dagger \subset A^\times$$

where $A^\dagger$ is the Hilbert space adjoint operator of A and $A^\dagger|_\Phi$ is its restriction to the subspace Φ . Generalized eigenvectors are defined for continuous operators. A vector $|F\rangle \in \Phi^\times$ is a generalized eigenvector of the $\mathcal{T}_\Phi$ -continuous operator A if for some complex number ω and for all $\phi \in \Phi$,

$$(3.4) \quad \langle A\phi|F\rangle = \langle \phi|A^\times|F\rangle = \omega \langle \phi|F\rangle$$

This is also written as

$$(3.5) \quad A^\times|F\rangle = \omega|F\rangle$$

(or, even as $A|F\rangle = \omega|F\rangle$ if $A^\dagger$ is a self-adjoint operator).

If A is the (self adjoint) Hamiltonian H of a quantum physical system, then $\Phi^\times$ contains the Dirac kets

$$(3.6) \quad H^\times|E^-\rangle = E|E^-\rangle, \quad E \geq 0$$

$\Phi^\times$ can also contain generalized eigenvectors with complex eigenvalues, as e.g.,

$$(3.7) \quad H^\times|E_R - i\Gamma/2^-\rangle = (E_R - i\Gamma/2)|E_R - i\Gamma/2^-\rangle,$$

which we call Gamow vectors or Gamow kets [22].

There is not only one space Φ , but there are many (locally convex, nuclear, countably normed) topologies $\mathcal{T}_\Phi$, which lead to different completions Φ of Φ_{alg} (with the same $\mathcal{H}$). The choice of Φ depends on the particular physical problem at hand, e.g., Φ can be chosen such that the algebra of observables of a particular physical system is an algebra of $\mathcal{T}_\Phi$ continuous operators.

Further, in section 2 we said that we need to distinguish meticulously between states and observables. In order to be able to also distinguish mathematically between states and observables we have to introduce one space for states, which we call Φ_- , and another space for observables, which we call Φ_+ . In general $\Phi_+ \neq \Phi_-$, but $\Phi_+ \cap \Phi_- \neq \{0\}$. The state prepared by the preparation apparatus (e.g., accelerator) we denote by ϕ^+ , thus $\phi^+ \in \Phi_-$. The observable registered by the registration apparatus (e.g., detector) we denote by $|\psi^-\rangle\langle\psi^-|$, thus $\psi^- \in \Phi_+$, (cf. Appendix B for the scattering experiment). Therefore we need two Rigged Hilbert Spaces, one for prepared in-states ϕ^+ :

$$(3.8) \quad \phi^+ \in \Phi_- \subset \mathcal{H} \subset \Phi_-^\times,$$

and the other for the registered observables $|\psi^-\rangle\langle\psi^-|$ or detected out-states ψ^- :

$$(3.9) \quad \psi^- \in \Phi_+ \subset \mathcal{H} \subset \Phi_+^\times$$

In here the space $\mathcal{H}$ is the same Hilbert space (with the same physical interpretation).

Mathematically one can define the spaces of the vectors Φ by the spaces of their energy wave functions $\langle E|\phi\rangle^1$:

$$(3.10) \quad \phi^+ \in \Phi_- \Leftrightarrow \langle^+ E|\phi^+\rangle \in \mathcal{S} \cap \mathcal{H}_-^2|_{\mathbb{R}^+}, \text{ (well behaved Hardy functions in } \mathcal{C}^-).$$

¹ In the same way as one can define the Hilbert space $\mathcal{H}$ by the space of Lebesgue square integrable functions $\mathcal{H} \ni h \Leftrightarrow h(E) \in L^2[0, \infty)$, where the functions $h(E)$ are uniquely determined only up to a set of Lebesgue measure zero, which is a complicated and unphysical notion, cf. section 3 ref. [26]

$$(3.11) \quad \psi^- \in \Phi_+ \Leftrightarrow \langle^- E | \psi^- \rangle \in \mathcal{S} \cap \mathcal{H}_+^2|_{\mathbb{R}^+}, \text{ (well behaved Hardy functions in } \mathcal{C}^+).$$

The notation in here is the following: $\mathcal{C}^+(\mathcal{C}^-)$ denotes the open upper (lower) half of the complex energy plane of the second Riemann sheet for the analytically continued S -matrix, and $\mathcal{H}_\mp^2$ denotes the Hardy class functions [23] and $\mathcal{S}$ the Schwartz space functions. This explains the notation Φ_- and Φ_+ for the spaces. The subscript refers to the subscript in the standard notation of mathematics for Hardy class functions ($\mathcal{H}_-^p, \mathcal{H}_+^p$ respectively). The superscripts for ϕ^+ (in-states) and ψ^- (out-states) are the most common convention in scattering theory, cf. Appendix B.

Thus, in the physical interpretation, for each species of quantum physical system one has a pair of RHS's, (3.8) and (3.9). Whereas the “in-state” $\phi^+ \in \Phi_-$ describes the state that is physically defined by the preparation apparatus, the “out-state” $\psi^- \in \Phi_+$ describes the observable that is physically defined by the registration apparatus.

It is by this clear differentiation between the set of vectors $\{\phi^+\}$ which are admitted as in-states and the set of vectors $\{\psi^-\}$ which are admitted as out-observables that the RHS-theory differs from the usual scattering theory, where $\{\phi^+\} = \{\psi^-\} = \Phi \subset \mathcal{H}$ (cf. the asymptotic completeness condition according to which $\{\phi^+\} = \{\phi^-\} = \mathcal{H}$). According to (3.10) and (3.11), Φ_- and Φ_+ are different dense subspaces of the same Hilbert space $\mathcal{H}$ (which are both complete with respect to a stronger topology than $\mathcal{T}_\mathcal{H}$) with

$$(3.12) \quad \Phi_+ \cap \Phi_- \neq \{0\}, \text{ and } \Phi = \Phi_+ + \Phi_- \text{ is also dense in } \mathcal{H}.$$

After the RHS's (3.8) and (3.9) have been chosen to be the Hardy class spaces (3.10) and (3.11), the semigroup of section 2 turns up naturally from the mathematics. How one could empirically conjecture the RHS's of Hardy class will not be discussed here [24].

To obtain the semigroups we start with the unitary group of time evolutions in the Hilbert space $\mathcal{H}$.

$$(3.13) \quad U(t) = e^{iHt}, \quad U^\dagger(t) = e^{-iHt}$$

where $U^\dagger(t)$ denotes the Hilbert space adjoint of $U(t)$.²

We first turn to the RHS (3.9) and consider

$$(3.14) \quad U_+(t) \equiv U(t)|_{\Phi_+} \subset U(t), \quad \text{and } U(t)^\dagger \subset U_+^\times(t)$$

It can be shown that, as a consequence of the mathematical properties of Φ_+ , the restriction of $U(t)$ to Φ_+ , $U_+(t)$, is a $\mathcal{T}_{\Phi_+}$ -continuous operator only for $0 \leq t < \infty$. Therefore its conjugate operator $U_+^\times(t)$, which is an extension of the Hilbert space adjoint operator $U^\dagger(t)$, is well defined (by (3.2)) and continuous for $0 \leq t \leq \infty$ only. Thus in $\Phi_+^\times$ we have only the semigroup

$$(3.15) \quad U_+^\times(t) = (e^{iHt}|_{\Phi_+})^\times \equiv e_+^{-iH^\times t}, \quad 0 \leq t < \infty$$

The same considerations apply to the other RHS (3.8). One considers $U_-(t) \equiv U(t)|_{\Phi_-} \subset U(t)$, and its conjugate $U(t)^\dagger \subset U_-^\times(t)$, and proves mathematically that $U_-(t)$ is a $\mathcal{T}_{\Phi_-}$ -continuous operator only for $-\infty \leq t \leq 0$. Therefore $U_-^\times(t)$ is defined and continuous for $-\infty \leq t \leq 0$ only and one has in $\Phi_-^\times$ the semigroup

$$(3.16) \quad U_-^\times(t) = \left(e^{iHt}|_{\Phi_-}\right)^\times \equiv e_-^{-iH^\times t}, \quad -\infty < t \leq 0$$

Thus in the RHS (3.8) for the prepared states one has the semigroup (3.16) for times $t \leq t_0 = 0$, and in the RHS (3.9) for the registered observables one has the semigroup (3.15) for times $t \geq t_0 = 0$. Since $t = t_0 = 0$ is the time by which the state has been prepared and the registration of the observable can begin, this separation of the (mathematical) group (3.13) into the two semigroups (3.16) and (3.15) reflects the situation envisioned on empirical

² Note that in $\mathcal{H}$ the right hand side of (3.13) is not defined by the exponential series

$$I + \frac{iHt}{1!} + \frac{(iH)^2 t^2}{2!} + \frac{(iH)^3 t^3}{3!} + \dots$$

which only converges with respect to $\mathcal{T}_{\mathcal{H}}$ on a dense subspace of analytic vectors in $\mathcal{H}$, but by the Stone- von Neumann calculus.

grounds in section 2. The scattering (e.g., resonance scattering) process is separated into two parts, the preparation part dealing with the preparation of the state $\phi^+ \in \Phi$ and the registration part dealing with the registration of the observable (or detection of the out-state) $\psi^- \in \Phi_+$. The time $t = 0(t_0)$ is the time at which the preparation is completed and the registration can commence; the meaning of t_0 will be discussed in detail in section 4.

In addition to the vectors ϕ^+ and ψ^- defined by the apparatuses, there also are the vectors in $\Phi_\pm^\times$ which are outside of $\mathcal{H}$:

$$(3.17) \quad |E, \theta_p, \varphi_p^\mp\rangle \in \Phi_\pm^\times, \quad (\text{Dirac's scattering states}),$$

where (θ_p, φ_p) denotes the direction of momentum;

and the

$$(3.18) \quad \psi^G = |E_R \mp i\Gamma/2, j, j_3^\mp\rangle \in \Phi_\pm^\times \quad (\text{Gamow's resonance states})$$

with the property

$$(3.19) \quad H^\times |E_R - i\Gamma/2^-\rangle = (E_R - i\Gamma/2) |E_R - i\Gamma/2^-\rangle.$$

In the RHS theory Dirac kets and Gamow vectors are mathematically very similar. Both are generalized eigenvectors of self-adjoint Hamiltonians in the sense of (3.2) and are equally well defined, (though the choice of spaces Φ for which Gamow kets can be defined is smaller than for Dirac kets since the former also requires some analyticity properties as for Hardy class spaces Φ_+). Dirac kets and Gamow kets just differ in their eigenvalues; whereas Dirac's scattering state vectors in $\Phi_+^\times$ or $\Phi_-^\times$ have real (except for the $\pm i0$) eigenvalues corresponding to the scattering energies, Gamow kets have complex eigenvalues corresponding to the resonance pole of the S -matrix (see below).

The Gamow vectors $\psi^G = |E_R - i\Gamma/2\rangle \sqrt{2\pi\Gamma} \in \Phi_+^\times$ have a semigroup time evolution and obey an exponential law:

$$(3.20) \quad \psi^G(t) \equiv U_+^\times(t)\psi^G = e^{-iE_R t} e^{-\Gamma t/2} |E_R - i\Gamma/2^-\rangle, \quad t \geq 0$$

This is a formal consequence of applying the right hand side of (3.15) to ψ^G and using (3.19). But for the mathematical proof of (3.20), in particular of the semigroup character, the whole mathematical apparatus of the RHS of Hardy class is needed [21].

There are other Gamow vectors $\tilde{\psi}^G = |E_R + i\Gamma/2^+\rangle \sqrt{2\pi\Gamma} \in \Phi_-^\times$, and there is another semigroup (3.16), $e_-^{-iH^\times t}$ for $t \leq 0$ in $\Phi_- \subset \mathcal{H} \subset \Phi_-^\times$ with the asymmetric evolution

$$(3.21) \quad \tilde{\psi}^G(t) = e^{-iH^\times t} |E_R + i\Gamma/2^+\rangle = e^{-iE_R t} e^{\Gamma t/2} |E_R + i\Gamma/2^+\rangle, \quad t \leq 0$$

Gamow vectors have the following features:

1. They are derived as functionals of the resonance pole term at $z_R = E_R - i\Gamma/2$ (and at $z_R^* = E_R + i\Gamma/2$) in the second sheet of the analytically continued S -matrix [9, 25].
2. They have a Breit-Wigner energy distribution $|\langle^- E | \psi^G \rangle|^2 = \frac{\Gamma}{2\pi} \frac{1}{(E - E_R)^2 + (\frac{\Gamma}{2})^2} \rightarrow \delta(E - E_R)$ for $\frac{\Gamma}{E_R} \rightarrow 0$ which extends to negative energy values on the second sheet indicated in the representation

$$(3.22) \quad |\psi^G\rangle = i\sqrt{\frac{\Gamma}{2\pi}} \int_{-\infty_{II}}^{+\infty} dE \frac{|E^-\rangle}{E - (E_R - i\Gamma/2)}$$

by $-\infty_{II}$ [9].

3. The decay probability $\mathcal{P}(t) = \text{Tr}(\Lambda |\psi^G\rangle \langle \psi^G|)$ of $\psi^G(t)$, $t \geq 0$, into the final non-interacting decay products described by Λ can be calculated as a function of time, and from this the decay rate $R(t) = \frac{d\mathcal{P}(t)}{dt}$ is obtained by differentiation [26]. This leads to an exact Golden Rule (with the natural line width given by the Breit-Wigner) and the exponential decay law

$$(3.23) \quad R(t) = e^{-i\Gamma t} \Gamma_\Lambda \quad t \geq 0$$

where Γ_Λ is the partial width for the decay products Λ ($\Gamma_\Lambda = \text{branching ratio} \times \Gamma$). In the Born approximation ($\psi^G \rightarrow f^D$, an eigenvector of $H_0 = H - V$; $\Gamma/E_R \rightarrow 0$; $E_R \rightarrow E_0$) this exact Golden Rule goes into Fermi's Golden Rule No. 2 of Dirac.

4. The Gamow vectors ψ_i^G are members of a “complex” basis vector expansion [25]. In place of the well known Dirac basis system expansion (Nuclear Spectral Theorem of the RHS) given by

$$(3.24) \quad \phi^+ = \sum_n |E_n\rangle (E_n|\phi^+) + \int_0^{+\infty} dE |E^+\rangle \langle^+ E|\phi^+\rangle$$

(where the discrete sum is over bound states, which we henceforth ignore), every prepared state vector $\phi^+ \in \Phi_-$ can be expanded as

$$(3.25) \quad \phi^+ = \sum_{i=1}^N |\psi_i^G\rangle \langle\psi_i^G|\phi^+\rangle + \int_0^{-\infty_{II}} dE |E^+\rangle \langle^+ E|\phi^+\rangle$$

(where $-\infty_{II}$ indicates that the integration along the negative real axis or other contour including around cuts is in the second Riemann sheet of the S -matrix). N is the number of resonances in the system (partial wave), each one occurring at the pole position $z_{R_i} = E_{R_i} - i\Gamma_i/2$. This allows us to mathematically isolate the exponentially decaying states ψ_i^G .

The “complex” basis system expansion is rigorous. The Weisskopf–Wigner approximate methods are tantamount to omitting the background integral, i.e.,

$$(3.26) \quad \phi^+ \stackrel{\text{W}\equiv\text{W}}{=} \sum_{i=1}^N |\psi_i^G\rangle c_i$$

For instance, for the $K_L - K_S$ system with $N = 2$,

$$\phi^+ = \psi_S^G b_S + \psi_L^G b_L$$

The properties (3.18)–(3.25) are not independently postulated conditions for the Gamow vectors but derived from each other in the mathematical theory of the RHS. One can start for instance with the most widely accepted definition of the resonance by the pair of poles of the S -matrix (B.3) at $z_R = E_R \mp i\Gamma/2$ and associate to it the Lippmann-Schwinger-Dirac ket $|z_R^\mp\rangle$ obtained from analytic continuation in (B.3). Then one obtains the Breit-Wigner energy distribution (3.22) from the Hardy class property (3.11) and vice

versa. From (3.22), using (3.11)– in particular the property of the Schwartz space $\mathcal{S}$ – one derives (3.19) as generalized eigenvalue equation $\langle \psi^- | H^{\times n} | z_R^- \rangle = z_R^n \langle \psi^- | z_R^- \rangle$, not only for $n = 1$ but for all powers n . The generalized eigenvalue equation (3.20) is also derived from the representation (3.22) but only for $t \geq 0$ because of the Hardy class property (which in turn was needed to justify the Breit-Wigner energy distribution for the pole term of the S-matrix). The Dirac basis vector expansion (3.24) is fulfilled for every RHS, e.g., when Φ is realized just by $\mathcal{S}$. The basis vector expansion (3.25) follows by analytic continuation and therefore requires the Hardy class property (3.10)(3.11). The derivation of the exact Golden Rule [26] uses in addition the Lippmann-Schwinger equation (B.4).

4. The Physical Interpretation and the Meaning of the Initial Time

The semigroup time evolution introduces a new concept — the time $t_0(=0)$ at which the preparation of the state is completed and the registration of the observable can begin. This is the most difficult new concept, because one is unprepared for it by the school of thinking based on the old time symmetric quantum mechanics. For the state of our universe as a whole, considered as a *closed* quantum mechanical system, there is no problem, because we deal only with one single system and the time t_0 is the time of the creation of this single universe (big bag time). Alternatively, we could consider this universe as a member of an ensemble of universes, of which we have access to only our universe. Then the probabilities (2.19a)- (2.26a) are the statistical probabilities (“relative frequencies”) of this ensemble and we have the usual interpretation of quantum mechanics, where the density operator ρ , W , the state vector ϕ^+ or wave function $\langle^+E|\phi^+\rangle$ is the mathematical representative of an ensemble of microsystems.

For an experiment performed on a quantum system in the laboratory, the states prepared by a macroscopic preparation apparatus, i.e., states described by $\phi^+ \in \Phi_-$ or $W = \sum_i w_i |\phi_i^+\rangle \langle \phi_i^+|$, $\phi_i^+ \in \Phi_-$, are best interpreted as ensembles (e.g., the proton or electron beam prepared by an accelerator). But there are other “states” which are prepared by a macroscopic apparatus in conjunction with a quantum scattering process (e.g., resonance scattering), which are best interpreted as states of a single microsystem. For their description the RHS offers, e.g., the Dirac kets (3.17) or the Gamow kets (3.18). From the basis vector expansion (3.25), we know that, mathematically, the apparatus-prepared state ϕ^+ can be represented as the sum of a Gamow vector ψ^G and a background integral. We shall now argue that the Gamow state can also be isolated experimentally and discuss its creation time t_0 and its asymmetric development in time.¹ This microphysical irreversibility is the analogue of the arrow of time for the state of our universe.

¹ For another discussion of the impossibility of time reversing the development of a decaying microphysical system, see T. D. Lee [16].

The best example is the decaying state of the neutral Kaon system because it is a wonderfully closed system, isolated from most external influences (including the electromagnetic field) whose (exact) evolution in time is probably entirely due to the Hamiltonian of the neutral Kaon system and free of external influences like those mentioned in footnote 1 of section 2. Since we are here only interested in the fundamental concepts of decay, we discuss a simplified K° -system for which the K_L° as well as the CP violation is ignored [27].

The process (idealized, because in the real experiment one does not use a π —but a proton beam) by which the neutral Kaon state is prepared is:

$$(4.1) \quad \pi^- p \Longrightarrow \Lambda K^\circ; \quad K^\circ \Longrightarrow \pi^+ \pi^-.$$

K° is strongly produced with a time scale of 10^{-23} sec. and it decays weakly, with a time scale of 10^{-10} sec, which is roughly the lifetime of the K_S° , τ_{K_S} . Thus t_0 , the time at which the preparation of the K° -state, which we call W^{K° , is completed and the registration can begin, is very well defined. (Theoretical uncertainty is $10^{-13}\tau_K$). A schematic diagram of a real experiment [28] is shown in Figure 1. The state W^{K° is created instantly at the baryon target T (and the baryon B is excited from the ground state (proton) into the Λ state, with which we are no further concerned). We imagine that a single particle K° is moving into the forward beam direction, because somewhere at a distance, say at d_2 from T , we “see” a decay vertex for $\pi^+\pi^-$, i.e., a detector (registration apparatus) has been built such that it counts $\pi^+\pi^-$ pairs which are coming from the position d_2 . The observable registered by the detector is the projection operator

$$(4.2) \quad \Lambda(t_2) = |\pi^+\pi^-, t_2\rangle\langle\pi^+\pi^-, t_2| = |\psi^{\text{out}}(t_2)\rangle\langle\psi^{\text{out}}(t_2)|$$

for those $\pi^+\pi^-$ which originate from the fairly well specified location d_2 . From the position (in the lab frame) d_2^{lab} , the four-momentum p of the K° (= the z component of the momentum of the $\pi^+\pi^-$ system) and the mass m_K of K° , one obtains the time t_2^{rest} (in

the K° rest frame) which the K° has taken to move from T to d_2^{lab} . This is given by the simple formula of relativity $d_2^{\text{lab}} = t_2^{\text{rest}} \frac{p}{m_K}$ which we write $d_2 = t_2 \frac{p}{m_K}$.

We do not have to focus at only one location d_2 but can count decay vertices at any distance d (of the right order of magnitude). The detector (described by the projection operator $\Lambda(t) \equiv |\pi^+\pi^-, t\rangle\langle\pi^+\pi^-, t|$) counts the $\pi^+\pi^-$ decays at different times $t = t_1, t_2, t_3, \dots$ (in the rest frame of the K°), and these correspond to the distances from the target $d_1 = pt_1/m_K, d_2 = pt_2/m_K, \dots$ (in the lab frame).

One “sees” the decay vertex d_i for each single decay and imagines a single decaying K° micro-system that had been created on the target T at time $t_0 = 0$ and then traveled a time t_i until it decayed at the vertex d_i . We give the following interpretation to these observations: a single microphysical decaying system K° described by W^{K° has been produced by a macroscopic registration apparatus and a quantum scattering process, at a time $t = 0$. Each count of the detector is the result of the decay of such a single microsystem. This particular microsystem has lived for a time t_i —the time that it took the decaying system to travel from the scattering center T to the decay vertex d_i . The whole detector registers the counting rate $\frac{\Delta N(t)}{\Delta t} \approx NR(t)$ as a function of d_i , i.e., of $t_i = \frac{m_K}{p}d_i$, for $\dots t_i > \dots t_2 > t_1 > t_0 = 0$. (N is the total number of counts).

The counting rate $\frac{\Delta N(t_i)}{\Delta t}$ is plotted as a function of time t (in the K° rest frame), in Figure 2.

No $\pi^+\pi^-$ are registered for $t < 0$, i.e., clicks of the counter for $\pi^+\pi^-$ that would point to a decay vertex at the position d_{-1} in front of the target T are not obtained (if there were any, they would be discarded as noise). One finds for the counting rate

$$(4.3) \quad \frac{\Delta N(t_i)}{\Delta t} \approx 0, \quad t < 0$$

This is so obvious that one usually does not mention it. For $t > 0$ one can fit the experimental counting rate with the exponential function to as good accuracy as one wants (by

taking larger N and smaller Δt):

$$(4.4) \quad \frac{\Delta N(t_i)}{\Delta t} \approx N e^{-\Gamma t} \quad t > 0 = t_0$$

The $\approx$ in (4.4) means, as in (2.1), the equality between experimental numbers and the idealized, theoretical hypothesis $e^{-\Gamma t}$ [29].

Theoretically, the counting rate is given by the probability rate

$$(4.5) \quad R(t) = \frac{d\mathcal{P}(t)}{dt}$$

where $\mathcal{P}(t)$ is the probability for the observable $\Lambda(t)$ of (4.2) (i.e., $\pi^+\pi^-$) in the state W^{K° .

According to the postulate (2.17), the probability should be given by,

$$(4.6) \quad \mathcal{P}(t) = \text{Tr}(\Lambda(t)W^{K^\circ}) = \text{Tr}(\Lambda W^{K^\circ}(t)) \text{ for } t \geq t_0 = 0$$

$$(4.7) \quad \text{where} \quad W^{K^\circ}(t) = e^{-iHt}W^{K^\circ}e^{iHt} \quad \text{for } t \geq t_0$$

For $t < t_0 = 0$, $W^{K^\circ}(t)$ is nonexistent because the K° had not been prepared by $t < t_0$.

To calculate theoretical results that agree with the observations (4.3) and (4.4), one has to choose the state operator $W^{K^\circ}(t)$ in (4.6) such that $W^{K^\circ}(t)$ is nonexistent for $t < t_0 = 0$, and such that for $t > t_0 = 0$, yields by (4.5) and (4.6), a result that is in agreement with the right hand side of (4.4). The state operator which has this property is given by (2.16),

$$(4.8) \quad W^{K^\circ}(t) = |F(t)\rangle\langle F(t)|$$

where

$$(4.9) \quad F(t) = U_+^\times(t)F_0$$

is a semigroup solution (2.15) of the quantum mechanical Cauchy problem, and where the initial vector is given by the Gamow vector

$$(4.10) \quad F_0 = |E_R - i\Gamma/2^-\rangle \in \Phi_+^\times$$

with $E_R = m_S$ and $\Gamma = \frac{1}{\tau_S}$ for the K° at rest [30].

Then we obtain the time evolved state vector (3.20) by applying the semigroup (3.15). For this vector $|E_R - i\Gamma/2^-\rangle$ (and only for this Gamow vector $|E_R - i\Gamma/2^-\rangle \in \Phi_+^\times$, which is defined by the pole term of the S -matrix) one derives the exact Golden Rule with the (exact) exponential decay law (3.23), thus reproducing the right hand side of (4.4).

Therewith we see that the Gamow state vector $\psi^G = |E_R - i\frac{\Gamma}{2}^-\rangle$ or the operator $W^G = |\psi^G\rangle\langle\psi^G|$, whose time evolution is governed by the exact Hamiltonian H , describes the decaying neutral Kaon system (4.1) in its rest frame if $E_R = m_s$ and $\Gamma = \Gamma_s = \frac{1}{\tau_s}$, $W^G = W^{K^\circ}$.

For this Gamow state one can calculate the decay rate and decay probability as a function of time and obtain the exponential law for $t > t_0 = 0$. The decay probability is the a priori probability for a single decaying microsystem K° that has been created in the state W^{K° at the initial time $t = 0$ (for the quantum mechanical Cauchy problem with semigroup evolution). This is the same point of view mentioned at the beginning of this section for the quantum state of our universe [3], except that its initial state $\rho(t_0)$ is probably not a “pure” Gamow state. Alternatively $W^{K^\circ}(t)$ can also be thought of as describing the state of an ensemble of single microsystems K° created at an “ensemble” of times t_0 , all of which are chosen to be the initial time $t = 0$ for the quantum mechanical Cauchy problem. Then the decay probabilities are the statistical probabilities for this ensemble of individual K° systems, but t in $W^{K^\circ}(t)$ is the time in the “life” of each single decaying K° -system which had started at $t = 0$. It is not the time in the experimentalists life or the time in the laboratory or the time of a “wave-packet” of K° ’s.

With this interpretation the single quasi-stable particle and the single quantum universe are perfectly analogous, and the time t_0 , at which the preparation of the state is completed and at which the registration of the observables can begin, has been observationally defined.

5. Summary and Conclusion

If we want to have a quantum theory that applies to the (closed) universe as a whole then we would like this quantum theory to be time asymmetric, because of the cosmological arrow of time. By the same reasoning if a quantum theory is to apply to the electromagnetic field then it should be time asymmetric, because of the radiation arrow of time. Standard quantum theory is time symmetric. This is a mathematical consequence due to the property of the Hilbert space postulates.

There is a mathematical theory that describes time symmetric as well as time asymmetric quantum physics. It is an extension of the Rigged Hilbert Space (RHS) formulation of quantum mechanics which about 1965 gave a mathematical justification to Dirac's kets and his continuous basis vector expansion.

To incorporate causality, the RHS theory distinguishes meticulously between states and observables for which it uses two RHS's $\Phi_{\mp} \subset \mathcal{H} \subset \Phi_{\mp}^{\times}$ of Hardy class with complementary analyticity properties. The dual spaces in the RHS's contain, besides the Dirac kets $|E^{\pm}\rangle \in \Phi_{\mp}^{\times}$ ($_{\text{out}}^{\text{in}}$ plane waves), also Gamow kets $|E_R - i\Gamma/2^{\pm}\rangle \in \Phi_{\mp}^{\times}$. The Gamow kets have all the properties attributed to eigenvectors of complex finite-dimensional Hamiltonian in the phenomenological effective theories, in particular an exponential time evolution and a Breit-Wigner energy wave function. Neither of this is possible in Hilbert space.¹ These effective finite dimensional theories can therefore not be considered approximations [15] of standard quantum mechanics, but they go beyond it. The RHS formulation is the mathematical theory of which these finite dimensional models, e.g., the two dimensional Lee-Oehme-Yang theory for neutral Kaons and the Weisskopf-Wigner method, are approximations.

Features of the exact theory which are not already features of these effective models and phenomenological methods are : the Breit-Wigner wave function (3.22) of the Gamow ket which extends over $-\infty_{\text{II}} < E < +\infty$ (rather than the values $0 \leq E < \infty$), the

¹ The Breit-Wigner energy wave function would not be in the domain of the Hamiltonian.

background integral in (3.25), and the exact Golden Rule [26] for the decay probability $P(t)$ from which the decay rate $R(t)$ with exponential time dependence (3.23) is obtained by differentiation. Dirac's Golden Rule for the initial decay rate is the Born approximation at $t = 0$ of this exact rule for the decay rate $R(t)$. These are features which one may welcome or accept. The most surprising, unwanted and mostly rejected feature of the exact RHS theory is the semigroup time evolution (3.20) and (3.21) of the Gamow state, which is a manifestation of a fundamental quantum mechanical arrow of time.

Decaying Gamow states can be experimentally isolated as quasistationary microphysical systems if their time of preparation can be accurately identified. The observed decay probabilities (4.3), (4.4) of the neutral Kaon system have the same features as derived from a Gamow state, including the time ordering, (4.6). This time ordering is the same as the postulated time ordering in the probabilities of the histories of the universe considered as a quantum system, (2.19)-(2.25). Under this hypothesis the fundamental quantum arrow of time – expressing the vague notion of causality – can be considered subsumed under the cosmological arrow of time.

Acknowledgement

The author would like to acknowledge the very valuable conversations on many points discussed in this paper with M. Gell-Mann, which were actually the origin of this article. He would also like to thank J. Hartle for clarifying remarks on their papers. S. Wickramasekara and H. Kaldass provided suggestions and help with the preparation of this article. Support from the Welch Foundation is gratefully acknowledged.

Appendix A

In the Hilbert space formulation of quantum mechanics, the linear scalar product space Φ_{alg} is completed with respect to the norm to obtain a Hilbert space $\mathcal{H}$. The Hamiltonian H in the Schroedinger–von Neumann equations (2.8) is self-adjoint and semi-bounded, and the initial data $\phi_0, \psi_0 \in \mathcal{H}$.

Then one has the following mathematical theorems:

1. (Gleason) For every probability $\mathcal{P}(\Lambda)$, there exists a positive trace class operator ρ such that $\mathcal{P}(\Lambda) = \text{Tr}(\Lambda\rho)$ [31].
2. (Stone–von Neumann) The solutions of the Schroedinger–von Neumann equations for this ρ are time symmetric and given by the group $U^\dagger(t) = e^{-iHt}$ of unitary operators [32].
3. (Hegerfeldt) For every Hamiltonian H (self-adjoint, semi-bounded),

either

$$\text{Tr}(\Lambda(t)\rho) = \text{Tr}(\Lambda\rho(t)) = 0 \quad \text{for } -\infty < t < \infty$$

or,

$$\text{Tr}(\Lambda(t)\rho) = \text{Tr}(\Lambda\rho(t)) > 0 \quad \text{for all } t \text{ (except on a set of Lebesgue measure zero)}.$$

Here, Λ can be any positive, self-adjoint operator such as $\Lambda = |\psi\rangle\langle\psi|$ or $\Lambda = C_\alpha$ of (2.20) and ρ any trace class operator like $\rho = |\phi\rangle\langle\phi|$ or $\rho = \rho(t_{\text{big bang}})$ [33].

Theorem 1 says that all probabilities must be given by the trace. From theorem 3 it then follows that there cannot be a state ρ in the Hilbert space $\mathcal{H}$ that has been created or prepared a finite time $t - t_0$ ago, and for which therefore $\text{Tr}(\Lambda\rho(t)) = 0$ for $t < t_0$, which for $t \geq t_0$ decays into decay products Λ with a decay probability that is different from zero. This means, there exist no elements ϕ in $\mathcal{H}$ that can represent decaying states. Also absent are the states $\rho^{\text{eff}}(t_i)$ that have been created at times $t = t_0 = t_{\text{big bang}}$, $t = t_1 > t_0$, $t = t_2 > t_1$, etc., and whose probabilities $\text{Tr}(P(t_n)\rho^{\text{eff}}(t_{n-1}))$ are different from zero. Theorem 2 prohibits the asymmetric time evolution of a state in $\mathcal{H}$ and therewith the existence of a distinguished time t_0 of creation.

Appendix B: S Matrix and Lippmann-Schwinger Equation

Every experiment consists of a preparation stage and a registration stage. In the preparation stage of the scattering experiment, a (mixture of) initial states ϕ^{in} is prepared before the interaction $V = H - K$ is effective (e.g., by an accelerator outside the interaction region of the target). The initial state vectors ϕ^{in} , describing the non-interacting beam and target, evolve in time according to the free Hamiltonian K : $\phi^{\text{in}}(t) = e^{-iKt}\phi^{\text{in}}$. When the beam reaches the interaction region, the free in-state ϕ^{in} turns into the exact state vector ϕ^+ whose time evolution is governed by the exact Hamiltonian $H = K + V$

$$(B.1) \quad \Omega^+\phi^{\text{in}}(t) \equiv \phi^+(t) = e^{-iHt}\phi^+ = \Omega^-\phi^{\text{out}}$$

This vector leaves the interaction region and ends up as the well determined state ϕ^{out} . The state vector ϕ^{out} is determined from ϕ^{in} by the dynamics of the scattering process:

$$(B.2) \quad \phi^{\text{out}} = S\phi^{\text{in}} \quad S = \Omega^{-\dagger}\Omega^+$$

ϕ^{in} is controlled and determined by the preparation apparatus. ϕ^{out} is also controlled by the preparation apparatus and is in addition determined by the interaction V .

In the registration stage, the detector outside the interaction region does not detect ϕ^{out} , but rather it detects an observable $\psi^{\text{out}}(t) = e^{iKt}\psi^{\text{out}}$ (or a mixture thereof). ψ^{out} is controlled by the registration apparatus (trigger, energy efficiency, etc., of the detector). The detector counts are a measure of the probability to find the observable (property) $|\psi^{\text{out}}\rangle\langle\psi^{\text{out}}|$ in the state ϕ^{out} . This probability $|(\psi^{\text{out}}, \phi^{\text{out}})|^2$ is calculated by the S -matrix.

The S -matrix is the probability amplitude $(\psi^{\text{out}}, \phi^{\text{out}})$ which is calculated in the following way:

$$(B.3) \quad \begin{aligned} (\psi^{\text{out}}, \phi^{\text{out}}) &= (\psi^{\text{out}}, S\phi^{\text{in}}) = (\Omega^-\psi^{\text{out}}, \Omega^+\phi^{\text{in}}) \\ &= (\psi^-, \phi^+) = \int_0^\infty dE \langle\psi^-|E^-\rangle S(E+i0) \langle^+E|\phi^+\rangle \end{aligned}$$

$\phi^+(t) = e^{iHt/\hbar}\phi^+$ comes from the prepared in-state $\phi^{\text{in}}(t \rightarrow -\infty) = (\Omega^+)^{-1}\phi^+(t \rightarrow -\infty)$.

The free observable vector ψ^{out} emerges from the observable vector ψ^- whose time evolution is governed by the exact Hamiltonian H .

$\psi^-(t) = e^{iHt/\hbar}\psi^-$ goes into the measured out-state $\psi^{out}(t \rightarrow +\infty) = (\Omega^-)^{-1}\psi^-(t \rightarrow +\infty)$. Ω^+ and Ω^- are the Møller wave operators. The Lippmann-Schwinger equation relates the (known) eigenvectors of the free Hamiltonian K to two sets of eigenvectors of the exact Hamiltonian H

$$(B.4) \quad \begin{aligned} |E^\pm\rangle &= |E\rangle + \frac{1}{E - K \pm i\epsilon} V |E^\pm\rangle \\ &= |E\rangle + \frac{1}{E - H \pm i\epsilon} V |E\rangle = \Omega^\pm |E\rangle \end{aligned}$$

where

$$(B.5) \quad K|E\rangle = E|E\rangle \quad , \quad H|E^\pm\rangle = E|E^\pm\rangle$$

This defines the exact energy wavefunctions in terms of the in- and out- energy wave functions, whose modulus gives the energy resolution of the experimental apparatuses:

$$(B.6) \quad \langle^+ E | \phi^+ \rangle = \langle E | \phi^{in} \rangle \text{ is the incident beam resolution, it describes the energy}$$

distribution given by the accelerator (preparation apparatus).

$$(B.7) \quad \langle^- E | \phi^- \rangle = \langle E | \phi^{out} \rangle \text{ is the energy distribution of the detected state, it is given by}$$

the energy resolution of the detector (registration apparatus).

Since ϕ^{in} is controlled by the preparation apparatus, so is ϕ^+ . Likewise, since ψ^{out} is controlled by the registration apparatus, so is ψ^- . All this is quite standard, cf. [34] chapter 7, except that of the two versions, mentioned on p.188 of [34] as equally valid descriptions, we allow only the first version which is in agreement with our physical intuition of causality. In order to do this we distinguish between the set of in-state vectors $\{\phi^+\} \equiv \Phi_-$ and the set of out-observable vectors $\{\psi^-\} \equiv \Phi_+$. This hypothesis is quite natural since the state ϕ^+ (or ϕ^{in}) must be prepared before the observable $|\psi^-\rangle\langle\psi^-|$ (or ψ^{out}) can be measured in it. As shall be discussed in section 3, Φ_- and Φ_+ are different dense subspaces of the same Hilbert space $\mathcal{H}$.

Footnotes and References

- [1] The literature on resonances and decay is so large that it is difficult to list here even a representative selection. The standard monograph is M. L. Goldberger, K. M. Watson, *Collision Theory*, Wiley, New York (1964). The irreversible character of quantum mechanical decay has rarely even been mentioned (exceptions are C. Cohen-Tannoudji, et al., *Quantum Mechanics*, Vol. II, p. 1345; Wiley, New York (1977); T. D. Lee, [16] below) and to our knowledge not been incorporated in a theory of decay.
- [2] We are *not* concerned here with irreversibility in the quantum theory of open systems for which the asymmetric time evolution is described by a Liouville equation containing terms for the effects of the external reservoir. Cf. footnote 1 of section 2.
- [3] M. Gell-Mann, J. B. Hartle, p. 311 in *Physical Origins of Time Asymmetry*, J. J. Halliwell, et al., eds., (Cambridge University, 1994).
- [4] B. A. Lippmann, J. Schwinger, *Phys. Rev.*, **79**, 469 (1950);
M. Gell-Mann, H. L. Goldberger, *Phys. Rev.*, **91**, 398 (1953).
- [5] R. Ritz, *Physikalische Zeitschrift*, **9** (1908) 903; R. Ritz, *Physikalische Zeitschrift*, **10** (1909) 224; R. Ritz and A. Einstein, *Physikalische Zeitschrift*, **10** (1909) 323.
- [6] J. von Neumann, *Mathematische Grundlagen der Quantentheorie*, (Springer, Berlin, 1931) (English translation. Princeton University Press, Princeton, 1955).
- [7] P. A. M. Dirac, *The Principles of Quantum Mechanics*, (Clarendon Press, Oxford, 1930).
- [8] E. Roberts, *J. Math. Phys.*, **7** (1966) 1097; A. Bohm, *Boulder Lectures in Theoretical Physics 1966*, Vol. 9A, (Gordon and Breach, New York, 1967); J. P. Antoine, *J. Math. Phys.*, **10** (1969) 53; **10** (1969) 2276; See also O. Melsheimer, *J. Math. Phys.*, **15** (1974) 902; 917.
- [9] A. Bohm, *J. Math. Phys.*, **22** (1981) 2813; *Lett. Math. Phys.* **3** 455 (1978).
- [10] Some (e.g. A. Pais, *CP Violation: the First 25 Years*, in *CP Violation*, J. Tran,

Thanh Van, Eds., Editions Frontiers, (1990)) reserve the name “particle” for an object with a unique lifetime (in addition to the unique mass), in distinction to such superpositions as $|K^\circ\rangle$ and $|\bar{K}^\circ\rangle$ which are the states in which the neutral Kaon system is prepared. In our theory, these exact prepared states like the $|K^\circ\rangle$ are the ϕ^+ of (3.25), which, in addition to the quasi-stable particle states like ψ_i^G representing K_i° , also contain a background integral.

- [11] V. Weisskopf and E. P. Wigner, Z. f. Physik, **63**, 54 (1930); **65**, 18 (1930); W. Heitler, *Quantum Theory of Radiation*, Oxford (1954).
- [12] T. D. Lee, R. Oehme and C. N. Yang, Phys. Rev., **106** (1957) 340.
- [13] *Non-diagonalizable* complex Hamiltonian matrices (Jordan blocks) led to Jordan vectors instead of eigenvectors. Jordan vectors have a non-exponential time evolution of the order of $\hbar/\Gamma$ (and are therefore ruled out as decaying vectors). H. Banmngärtel *Analytic Perturbation Theory for Matrices and Operators*. Operator theory. Vol. 15. Birkhäuser Basel (1985), chapter 2; P. V. Ruuskanen, Nucl. Phys. **B22**, 253 (1970); E. Katznelson, J. Math. Phys. **21**, 1393 (1980); E. Hernandez and A. Mondragon, Phys. Lett. **B326**, 1 (1994); A. Mondragon and E. Hernandez, J. Phys. **A26**, 5595 (1993). However, the Jordan block Hamiltonians can also be shown to result from a truncation of the exact Rigged Hilbert Space theory to an effective theory. Its higher order Gamow states, which correspond to higher order S -matrix poles, and are not representable by vectors but by state operators have an exponential asymmetric time evolution. A. Bohm, M. Loewe, S. Maxson, P. Patuleanu, C. Püntmann and M. Gadella, J. Math. Phys., **38**, 6072 (1997).
- [14] E. Maglione, L. S. Ferreira and R. J. Liotta, Phys. Rev. Lett., **81** (1998) 538; O. I. Tolstikhin, V. N. Ostrovsky and H. Nakamura, Phys. Rev. **A58**, 2077 (1998); H. Hogreve, Phys. Lett. **A201**, 111 (1995); L. S. Ferreira in *Resonances*, E. Brändas et al. [Eds] Lecture Notes in Physics Vol. **325** p.201, Springer Berlin (1989).
- [15] M. Levy, Nuovo Cimento, **13**, 115 (1959).

- [16] T. D. Lee, Chapter 13 of *Particle Physics and Introduction to Field Theory*, Harwood Academic, New York, 1981. In this reference the quantum mechanical time reversed state is called complicated and improbable.
- [17] A. Kossakowski, On Dynamical Semigroups and Open Systems, in *Irreversibility and Causality*, A. Bohm, H. D. Doebner, P. Kielanowski [Eds.] Springer, Berlin (1998), p. 59, where further references to this irreversibility of quantum statistics can be found.
- [18] G. Ludwig, *Foundations of Quantum Mechanics*, Volume I, Springer-Verlag, Berlin, (1983) and Volume II, (1985); *An Axiomatic Basis of Quantum Mechanics*, Volume I, Springer-Verlag, Berlin, (1983) and Volume II, (1987).
- [19] M. Gell-Mann and J. B. Hartle, in *Complexity, Entropy and the Physics of Information*, SFI Studies in Science and Complexity Vol. VIII, W. Zurek, Ed., (1990).
- [20] M. Gell-Mann, J. B. Hartle, *Proceedings, 4th Drexel Symposium on Quantum Non-Integrability*, preprint UCSBTH-95-28.
- [21] A. Bohm and M. Gadella, *Dirac Kets, Gamow Vectors and Gel'fand Triplets*, Lecture Notes in Physics, **348**, Springer-Verlag, Berlin, (1989).
- [22] The Gamow vectors were envisioned by G. Gamow (Z. Physik, **51**, 204 (1928)) as far back as the inception of the Dirac kets. Mathematically, they were no more ill defined than the Dirac kets, though they never acquired the popularity of the latter. The reason was that, although for Dirac kets at least the probability density is finite everywhere, the probability density of Gamow's wavefunctions $|\langle r, \theta, \varphi | \psi^G(t) \rangle|^2$ increased exponentially for large values of the distance r and large negative values of time t , (the "exponential catastrophe"). The origin of this exponential catastrophe is, that the emission of decay products had been assumed to go on for an arbitrarily long time, $t \rightarrow -\infty$, as dictated by the unitary group e^{-iHt} , $-\infty < t < \infty$. In reality, the emission of decay products must have begun at some finite time $t = t_0 = 0$ in the past, and the time evolution of the decaying state, as we now

know, is described by the semigroup $e^{-iH^\times t}$, $0 < t < \infty$. Therefore, in the time asymmetric theory the probability density $|\langle r, \theta, \varphi | \psi^G(t) \rangle|^2$ is only defined for times $t > t_0(r) = \frac{r}{v} = \frac{mr}{p} \approx \frac{mr}{\sqrt{2mE_R}}$, and for these times one can show that there exists no exponential catastrophe for the probability densities. Thus the semigroup evolution of the Gamow vectors is not only a consequence of the mathematical theory, but also a necessity for the physical interpretation of the position probabilities $\int_{\Delta\vec{r}} r^2 dr d\theta d\varphi |\langle r, \theta, \varphi | \psi^G(t) \rangle|^2$.

- [23] The function $G_+(E) (= \langle -E | \psi^- \rangle = \langle E | \psi^{\text{out}} \rangle)$ is a very well behaved function of the upper half plane $\mathcal{C}^+$ if it is well behaved, i.e. $G_+(E) \in \mathcal{S}$ (Schwartz space) and if it is the boundary value of an analytic function $G_+(z)$ in the upper half plane $\mathcal{C}^+$ which vanishes faster than any power at the infinite semicircle (i.e. $G_+(E) \in \mathcal{H}_+$). Similarly the function $G_-(E) (= \langle +E | \phi^+ \rangle = \langle E | \phi^{\text{in}} \rangle)$ is a very well behaved function of the lower half plane $\mathcal{C}^-$ if it is well behaved ($G_-(E) \in \mathcal{S}$) and if it is the boundary value of an analytic function in the lower half plane $\mathcal{C}^-$ which vanishes sufficiently fast at the lower semicircle ($G_-(E) \in \mathcal{H}_-$). For the definition of Hardy class functions and their mathematical properties needed here see Appendix A.2 of reference [25] and P. L. Duren, *$\mathcal{H}^p$ Spaces*, Academic Press, New York, (1970).
- [24] A. Bohm, I. Antoniou, P. Kielanowski, Phys. Lett. A **189** 442 (1994); A. Bohm, I. Antoniou, P. Kielanowski, J. Math. Phys., **36**, 2593 (1995).
- [25] A. Bohm, S. Maxson, M. Loewe, M. Gadella, Physica A **236**, 485 (1997).
- [26] A. Bohm, N. L. Harshmann in *Irreversibility and Causality*, p. 225, Sect. 7.4; A. Bohm, H. D. Doebner, P. Kielanowski [Eds.] Springer, Berlin (1998).
- [27] For a detailed analysis of the $K_L - K_S$ system in the framework of time asymmetric quantum mechanics with and without CP violation, see A. Bohm; hep-th/970542.
- [28] This is the simplified schematic diagram of several generations of experiments measuring CP violation in the neutral Kaon system; J. Christenson, J. Cronin, V. Fitch, R. Turley, Phys. Rev. Lett., **13**, 138 (1964); K. Kleinknecht, *CP Violation*, p. 41,

C. Jarskog (Ed.), World Scientific (1989) and references therein; NA**31**, G. D. Barr et al., Phys. Lett. B**317**, 233 (1993); E731; L. K. Gibbons, et al., Phys. Rev. Lett., **70**, 1199 and 1203 (1993).

- [29] This approximate agreement between a sequence of rational numbers $\frac{\Delta N(t)}{\Delta t}$ on the experimental side and a continuous function of real numbers $e^{-\Gamma t}$ on the theoretical side is the fundamental limit to which the exponential law can be verified; experimental limitations given by the resolution, e.g., the finite size of Δt , are still more important. Therefore any *infinitesimal* (not given in terms of the scale $\frac{1}{\Gamma}$) deviations from the exponential law that are derived from a mathematical theory (e.g., the Hilbert space idealization, L. A. Khalin, JETP Lett., **15**, 388 (1972)), are physically meaningless. More important for observed deviations from the exponential law, (e.g. S. R. Wilkinson et. al., Nature **387**, 575 (1997)), are the limitations on the preparation side of the experiment. According to (3.25) the prepared state ϕ^+ contains, in addition to the exponentially decaying Gamow state ψ^G (assuming $N = 1$), the background integral, which does not have exponential time evolution. Theoretically, the background term could be infinitesimally small, but since it depends upon the state preparation i.e., the function $\langle^+ E | \phi^+ \rangle = \phi^{\text{in}}(E)$, $0 \leq E < \infty$, the effect of the background term can be substantial. Experimental conditions may thus not allow the isolation of the Gamow state ψ^G from this background term, and consequently deviations from the exponential law will be observed. This is a familiar effect in resonance scattering experiments where deviations from the pure Breit-Wigner distribution of the Gamow state are observed and attributed to the background phase shifts, see e.g., A. Bohm, *Quantum Mechanics*, 3rd ed., sections XVIII.6–9, and XX.3, Springer-Verlag, New York, (1994). It is not experimental evidence against the existence of a Gamow state with Breit-Wigner energy distribution and exponential time evolution.

- [30] The K° is a relativistic decaying system and our discussion here is in terms of

the non-relativistic Gamow vectors. Relativistic Gamow kets can be defined from the poles of the relativistic S -matrix at the value of the invariant mass square $s_R = (m_R - i\frac{\Gamma}{2})^2$. They have at rest the same semigroup time evolution as (3.20) with $E_R \rightarrow m$ and $\Gamma \rightarrow \frac{\hbar}{\tau_R}$. A. Bohm, H. Kaldass, et al., (in preparation).

- [31] A. M. Gleason, J. Math., **6**, 885 (1957)
- [32] M. H. Stone, Ann. of Math. **33**, 643 (1932)
- [33] G. C. Hegerfeldt, Phys. Rev. Lett. **72**, 596 (1994)
- [34] R. G. Newton, *Scattering Theory of Waves and Particles* (McGraw-Hill, 1966)




Figure 1 Schematic diagram of the neutral K-meson decay experiment




Figure 2 K_S decay versus proper time